Catasetum

Kat-ah-SEE-tum

This unusual group of orchids offers fascinating, waxy flowers that often have the peculiar habit of discharging their pollen masses (pollinia) onto pollinators. Almost always deciduous, the pseudobulbous plants have strict growing and resting periods. Most flower before entering a dormant period when they drop their leaves.

**LIGHT** should be strong, especially near the end of the growth period. Early in the annual growth cycle, plants will tolerate less light, from 1,500 to 3,000 foot-candles. Plants grow best with light levels of 3,000 to 6,000 foot-candles, or one-half to three-fourths full sun. As pseudobulbs mature, harden them by giving slightly more light.

**TEMPERATURES** reflect the fact that these orchids are native to hot tropical areas and grow during the rainy summer months. During this growing period, day temperatures of 80 to 100 F and night temperatures of 60 to 65 F are beneficial. After growths mature, temperatures can be reduced to 55 F at night, with day temperatures of 70 to 85 F.


WATER is a critical factor for the production of large pseudobulbs that result in best flowering. A great quantity of water must be stored by the plant in a relatively short growing season. Water heavily as new leaves are forming. As the pseudobulb matures, gradually reduce watering frequency. Leaves will yellow and drop. At this time, watering should be stopped completely until new growth begins. Water lightly to rehydrate the pseudobulbs if shriveled severely.

HUMIDITY should be 40 to 60 percent. This can be provided in the home by placing the plants on trays of gravel, only partially filled with water so that the plants do not sit in the water. Air should always be moving around the plants to prevent fungal or bacterial disease, especially if high humidity or cool temperatures exist. In the greenhouse, the humidity is best

increased by use of a humidifier while cooling the air.

FERTILIZE and water regularly to produce strong pseudobulbs. Use a high nitrogen formulation (such as 30-10-10) while plants are in active growth, slowly tapering off as pseudobulbs forms. A blossombooster formulation (such as 10-30-20) should be used in the autumn, except for plants that normally bloom in the spring. Frequent applications of a dilute concentrations of fertilizer are more effective than occasional applications of strong concentrations.

POTTING should be timed to coincide with the initiation of new growth, usually in the spring. New roots will be produced quickly at that time, and plants will experience minimal setback. These plants have vigorous root systems and require a rich, moist potting medium during the growing season. Many growers bare-root their plants during the resting period to ensure dryness at that time. Fine-grade media are common for smaller pots; medium-grade media are used only in larger pots. Sphagnum moss is used successfully for plants in many areas, as it provides tremendous water and fertilizerholding capacities. Some plants can be grown on slabs of tree fern or other material, which makes it easier to keep them dry during dormancy; however, it is harder to keep them moist while growing. When well-grown, these orchids can be divided down to one mature pserdobulb and will then flower on the next mature growth.



The American Orchid Society is the world's leading provider of information about and related to orchids. We invite you to join us and learn about the world's most fascinating flowers and plants. Your membership entitles you to our monthly award-winning magazine, *Orchids*, 10 percent discount on items purchased at the AOS online store, free and discounted admission to more than 200 botanical gardens and arboreta and exclusive members only online content.

## **American Orchid Society**

At Fairchild Tropical Botanic Garden 10901 Old Cutler Road, Coral Gables, FL 33156 www.aos.org